arXiv:1005.4443v2 [quant-ph] 9 Mar 2011

Heisenberg-limited sensitivity with decoherence-enhanced measurements

[1,2]Daniel Braun and [1,2,3]John Martin
[1]*Université de Toulouse, UPS, Laboratoire de Physique Théorique (IRSAMC), F-31062 Toulouse, France*
[2]*CNRS, LPT (IRSAMC), F-31062 Toulouse, France and*
[3]*Institut de Physique Nucléaire, Atomique et de Spectroscopie, Université de Liège, 4000 Liège, Belgium*

Quantum-enhanced measurements use quantum mechanical effects in order to enhance the sensitivity of the measurement of classical quantities, such as the length of an optical cavity. The major goal is to beat the standard quantum limit (SQL), i.e. an uncertainty of order $1/\sqrt{N}$, where N is the number of quantum resources (e.g. the number of photons or atoms used), and to achieve a scaling $1/N$, known as the Heisenberg limit. So far very few experiments have demonstrated an improvement over the SQL. The required quantum states are generally highly entangled, difficult to produce, and very prone to decoherence. Here, we show that Heisenberg-limited measurements can be achieved without the use of entangled states by coupling the quantum resources to a common environment that can be measured at least in part. The method is robust under decoherence, and in fact the parameter dependence of collective decoherence itself can be used to reach a $1/N$ scaling.

Quantum mechanical noise imposes fundamental limitations on any measurement. The best–known example is the Heisenberg uncertainty relation, which provides a lower bound on the product of fluctuations of two non-commuting quantized variables. But even a classical system parameter x can in general not be measured with arbitrary precision with a finite number of measurements due to the statistical nature of any quantum state. A lower bound on the uncertainty is given by the smallest δx such that two quantum states $\rho(x)$ and $\rho(x+\delta x)$ lead to statistically significant differences for an optimally chosen observable. A similar problem exists in classical statistical analysis, where one wants to distinguish between two probability distributions $P(x)$ and $P(x+\delta x)$, and the celebrated Cramér-Rao bound sets an ultimate lower bound on the uncertainty of a measurement of x based on the distinguishability of $P(x)$ and $P(x+\delta x)$ [1]. That analysis has been generalized to the quantum world [2] and has become known as "quantum parameter estimation theory". Recently, the theory was used to prove lower bounds on the smallest measurable δx for unitary time evolution. It was shown that if N replicas of the quantum system evolve independently and linearly, for an initially separable state no uncertainty smaller than $1/\sqrt{N}$ can be achieved, i.e. the SQL, no matter how sophisticated the measurement. If initially entangled states are allowed, a $1/N$ scaling of the uncertainty is the ultimate lower bound under otherwise identical conditions [3–6]. The use of non-classical states of light for Heisenberg limited interferometry, notably the use of squeezed states, was proposed theoretically already in 1981 [7]. So-called NOON states have been investigated for super-resolution [8–10]. However, decoherence of these highly non-classical states has so far prevented reaching an uncertainty that scales as $1/N$ for systems with $N \gg 1$ [11, 12]. In [13] entanglement-free Heisenberg-limited sensitivity of a phase-shift measurement was reported for several hundred quantum resources by passing light many times through the phase-shifter.

Decoherence arises when a quantum system interacts with an environment with many uncontrolled degrees of freedom, such as the modes of the electromagnetic field, phonons in a solid, or simply a measurement instrument [14]. Decoherence destroys quantum mechanical coherence, and plays an important role in the transition from quantum to classical mechanics [15]. It becomes extremely fast for a mesoscopic or even macroscopic "distance" between the components of a "Schrödinger cat"-type superposition of quantum states. Universal power laws rule the scaling of the decoherence rates in this regime [16, 17]. Only recently could the collapse be time-resolved in experiments with relatively small "Schrödinger cat"–states [18, 19]. However, decoherence can depend very sensitively on the initial state and the coupling to the environment. Entire decoherence-free subspaces (DFS) can exist if the coupling operators to the environment have degenerate eigenvalues [20–24].

We show below that a collective coupling that depends on a parameter x of N quantum systems $\mathcal{S}_i$ to a common "environment" $\mathcal{R}$ can be used to measure x with an uncertainty that scales as $1/N$ with an initial product state of all subsystems. The method works whether $\mathcal{R}$ is entirely under our control, or a reservoir with many degrees of freedom to which we have only partly access, i.e. a collective decoherence process of the $\mathcal{S}_i$, as long as we can measure an observable of the environment.

Results

Model

Consider N quantum systems $\mathcal{S}_i$ coupled to a common environment $\mathcal{R}$. The hamiltonian of the total system has the form

$$H(x)=\sum_{i=1}^{N} H_i+\sum_{i,\nu} S_{i,\nu}(x)\otimes R_\nu+H_R\,, \tag{1}$$

where H_i is the hamiltonian of system $\mathcal{S}_i$, and for simplicity we take the $\mathcal{S}_i$ as non-interacting. H_R denotes the hamiltonian of $\mathcal{R}$, which may be itself a composite quantum system. Hamiltonian (1) can be a model of decoherence (in which case $\mathcal{R}$ would be the ensemble of many degrees of freedom of a "reservoir" to which we have only partial access), or $H(x)$ can generate a unitary evolution if $\mathcal{R}$ and $\mathcal{S}=\{\mathcal{S}_1,\ldots,\mathcal{S}_N\}$ are completely under our control. The sum over ν runs over an arbitrary number of operators for each subsystem $\mathcal{S}_i$ and $\mathcal{R}$, but R_ν can also mean operators on different subsystems of $\mathcal{R}$ if $\mathcal{R}$ is composite (e.g. positions of harmonic oscillators modelling a heat bath). In order to have a generic name for $\mathcal{R}$ that encompasses these different situations, we will refer to $\mathcal{R}$ as the "quantum bus". The entire dependence on x is included in the coupling operators $S_{i,\nu}(x)$. With "collective couplings" (and with "collective decoherence" if $\mathcal{R}$ is a reservoir) we mean $S_{i,\nu}(x)$ which do not depend on i.

The smallest uncertainty δx with which x can be measured is found from quantum parameter estimation theory [2]. If the state of a system is given by a density matrix $\rho(x)$, the smallest achievable δx is given by

$$\delta x\geq\delta x_{\min}=\frac{1}{\sqrt{M}\left(\frac{ds^2}{dx^2}\right)^{1/2}}\,, \tag{2}$$

where we allow for M repetitions of the same measurement in identically prepared states $\rho(x)$, and ds^2 is a metric on the space of density operators. It is related to the Bures' metric $d_{\text{Bures}}(\rho,\rho+d\rho)$, with $d\rho=\rho'(x)dx$, by $ds=2d_{\text{Bures}}(\rho,\rho+d\rho)$. For pure states, the Bures distance reduces essentially to their overlap, $d_{\text{Bures}}(|\psi\rangle\langle\psi|,|\phi\rangle\langle\phi|)=\sqrt{2}\sqrt{1-|\langle\psi|\phi\rangle|}$ [25]. If $\rho(x)$ and $\rho(x)+d\rho$ are related through a unitary transformation with generator $\hat{h}$, $\rho(x+dx)=\exp(-\mathrm{i}\hat{h}dx)\rho(x)\exp(\mathrm{i}\hat{h}dx)$, then [2]

$$d_{\text{Bures}}(\rho(x),\rho(x)+d\rho)=\langle\Delta\hat{h}^2\rangle^{1/2}dx\,. \tag{3}$$

An operational definition of δx is given by

$$\delta x=\frac{\langle\Delta A^2\rangle_x^{1/2}}{\sqrt{M}|\partial\langle A\rangle_x/\partial x|}\,, \tag{4}$$

where we see that δx corresponds to the quantum uncertainty of an observable A in state $\rho(x)$, suitably translated by the slope of $\langle A\rangle_x$ into a fluctuation of x. As usual, for any observable A, $\langle\Delta A^2\rangle\equiv\langle A^2\rangle-\langle A\rangle^2$, and all expectation values are with respect to $\rho(x)$. Inequality (2) holds for all possible measurements, and for $M\to\infty$ a measurement exists that saturates the bound [2].

Model (1) cannot be solved in all generality. However, if the interaction is sufficiently weak it can be treated in perturbation theory. Since we start in an initial product state at $t=0$, we can then relate properties of the full model to the single particle dynamics of all $\mathcal{S}_i$ and $\mathcal{R}$. We first establish the fundamental lower bound on δx with the help of quantum parameter estimation theory, and then calculate δx for a given measurement on $\mathcal{R}$.

Quantum parameter estimation theory

We decompose $H(x)=H_0+H_I(x)$, $H_I(x)=\sum_{i,\nu}S_{i,\nu}(x)\otimes R_\nu$, and switch to the interaction picture with respect to H_0, with wave function $|\psi_I(x,t)\rangle=\exp(\mathrm{i}H_0t)|\psi(x,t)\rangle$, $|\psi(x,t)\rangle=\exp(-\mathrm{i}H(x)t)|\psi_0\rangle$. In "Methods" we show that the Bures distance between the two states $\rho(x)=|\psi_I(x,t)\rangle\langle\psi_I(x,t)|$ and $\rho(x+dx)=|\psi_I(x+dx,t)\rangle\langle\psi_I(x+dx,t)|$ is given by

$$d^2_{\text{Bures}}(\rho(x),\rho(x+dx)) \;=\; dx^2\int_0^t\int_0^t dt_1dt_2K_{|\psi_0\rangle}(H_I'(x,t_1),H_I'(x,t_2)) \tag{5}$$

where we have defined the correlation function for any two operators A, B in the state $|\psi\rangle$, $K_{|\psi\rangle}(A,B)=\langle\psi|AB|\psi\rangle-\langle\psi|A|\psi\rangle\langle\psi|B|\psi\rangle$, $H_I(x,t)=\exp(\mathrm{i}H_0t)H_I(x)\exp(-\mathrm{i}H_0t)$ is the interaction hamiltonian in the interaction picture, and $H_I'(x,t)=\partial H_I(x,t)/\partial x$. Equation (5) generalizes (3), which is recovered if $[H'(x),H(x)]=0$ and $H_I(x)t=x\hat{h}$. From (2), we have

$$\delta x_{\min}=\frac{1}{2\sqrt{M}\left(\int_0^t\int_0^t dt_1dt_2K_{|\psi_0\rangle}(H_I'(x,t_1),H_I'(x,t_2))\right)^{1/2}}\,. \tag{6}$$

For identical and identically prepared systems $\mathcal{S}_i$, $S_{i,\nu}=S_\nu$ and $|\varphi\rangle_i=|\varphi\rangle$ for all i, and an initial product state

$$|\psi_0\rangle=|\varphi\rangle^{\otimes N}\otimes|\xi\rangle\,, \tag{7}$$

we find

$$\begin{aligned}K_{|\psi_0\rangle}(H_I'(x,t_1),H_I'(x,t_2)) &= \sum_{\nu,\mu}\Big(NK_{|\varphi\rangle}(S_\nu'(x,t_1),S_\mu'(x,t_2))\langle R_\nu(t_1)R_\mu(t_2)\rangle\\ &\quad+N^2\langle S_\nu'(x,t_1)\rangle\langle S_\mu'(x,t_2)\rangle K_{|\xi\rangle}(R_\nu(t_1),R_\mu(t_2))\Big)\,,\end{aligned} \tag{8}$$

where the expectation values for operators of $\mathcal{S}_i$ ($\mathcal{R}$) are taken in states $|\varphi\rangle$ ($|\xi\rangle$). Together with eq. (6) this proves the existence of a measurement on $\mathcal{S}$ and $\mathcal{R}$ that gives a $1/N$ scaling of $\delta x_{\min}$ for $N\gg1$ and an initial product state, provided $\sum_{\nu,\mu}\int_0^t\int_0^t\langle S_\nu'(x,t_1)\rangle\langle S_\mu'(x,t_2)\rangle K_{|\xi\rangle}(R_\nu(t_1),R_\mu(t_2))dt_1dt_2\neq0$.

Measuring the quantum bus

We now use directly eq. (4) for showing that the $1/N$ scaling can be achieved with the measurement of almost any observable A on $\mathcal{R}$ alone. The expectation values in eq. (4) are in general time-dependent. This implies a time-dependent minimal uncertainty as well which does, however, not affect the scaling with N. We evaluate $\langle A(t)\rangle$ and $\langle\Delta A^2(t)\rangle$ again by using second order perturbation theory in the interaction. The general results for these expressions are cumbersome, but simplify considerably if we make the following two assumptions: (1) The initial state of $\mathcal{R}$ is an eigenstate of A, $A|\xi\rangle=a_\xi|\xi\rangle$; and (2) A commutes with H_R. Both assumptions taken together imply that the quantum bus is prepared in a noiseless state at $t=0$ ($\langle\Delta A^2(0)\rangle=0$). Under the above two assumptions, we find

$$\begin{aligned}\langle A(t)\rangle &= a_\xi+\int_0^t dt_1\int_0^t dt_2\chi_{S\nu\mu}(N,x,t_1,t_2)C_{RA\nu\mu}(t_1,t_2)+\mathcal{O}(H_I^3)\,, && (9)\\ \chi_{S\nu\mu}(N,x,t_1,t_2) &= NK_{|\varphi\rangle}(S_\nu(x,t_1),S_\mu(x,t_2))+N^2\langle S_\nu(x,t_1)\rangle\langle S_\mu(x,t_2)\rangle && (10)\\ C_{RA\nu\mu}(t_1,t_2) &= \langle R_\nu(t_1)AR_\mu(t_2)\rangle-a_\xi\langle R_\nu(t_1)R_\mu(t_2)\rangle\,, && (11)\end{aligned}$$

which can also be used to obtain $\langle A^2(t)\rangle$ and $\langle\Delta A^2(t)\rangle$. Eq.(4) then leads to

$$\delta x=\frac{\left[\int_0^t dt_1\int_0^t dt_2\sum_{\nu,\mu}\chi_{S\nu\mu}(N,x,t_1,t_2)\langle[R_\nu(t_1),A][A,R_\mu(t_2)]\rangle\right]^{1/2}}{\sqrt{M}\left|\int_0^t dt_1\int_0^t dt_2\sum_{\nu,\mu}\frac{\partial}{\partial x}\chi_{S\nu\mu}(N,x,t_1,t_2)\langle R_\nu(t_1)[A,R_\mu(t_2)]\rangle\right|}\,. \tag{12}$$

In the limit of $N\gg1$, the term quadratic in N in $\chi_{S\nu\mu}(N,x,t_1,t_2)$ dominates, and we find a $1/N$ scaling of δx,

$$\delta x=\frac{\left[\int_0^t dt_1\int_0^t dt_2\sum_{\nu,\mu}\langle S_\nu(x,t_1)\rangle\langle S_\mu(x,t_2)\rangle\langle[R_\nu(t_1),A][A,R_\mu(t_2)]\rangle\right]^{1/2}}{\left|\int_0^t dt_1\int_0^t dt_2\sum_{\nu,\mu}\frac{\partial}{\partial x}\left(\langle S_\nu(x,t_1)\rangle\langle S_\mu(x,t_2)\rangle\right)\langle R_\nu(t_1)[A,R_\mu(t_2)]\rangle\right|}\frac{1}{\sqrt{M}N}\,, \tag{13}$$

provided that the denominator does not vanish. It is enough to measure an observable of the quantum bus $\mathcal{R}$ alone, with all subsystems initially in a product state.

Decoherence

All derivations so far apply perfectly well if $\mathcal{R}$ is an environment with many degrees of freedom which we cannot fully measure. Measuring an observable A on only a subset of these implies a non-unitary evolution of $\mathcal{S}$. This establishes immediately that we can reach a $1/N$ scaling of δx, if x parametrizes a collective decoherence process, and if we can measure at least some part of the environment. The example of superradiance that we will work out below is of this type. However, one might also be interested in how the unitary evolution generated by the hamiltonian (1) is affected by additional independent decoherence of the components $\mathcal{S}_i$ and $\mathcal{R}$. For Markovian decoherence, such a situation is described by a master equation for the density matrix $W(t)$ of $\mathcal{S}$ and $\mathcal{R}$ of the form

$$\dot{W}(t) = -\mathrm{i}\left(L_0 + L_I(x) + \mathrm{i}\Lambda_s + \mathrm{i}\Lambda_R\right) W(t), \tag{14}$$

where $L_0\, X = [H_0, X]$, $L_I(x)\, X = [H_I(x), X]$, and Λ_R (Λ_S) are Liouvillians of the Lindblad-Kossakowski type [26] for $\mathcal{R}$ ($\mathcal{S}$), with $\Lambda_S = \sum_{i=1}^{N} \Lambda_i$.

The free evolution ($H_I = 0$) still factorizes, such that, essentially, all expectation values and correlation functions are replaced by expectation values with respect to the relevant mixed states (see eq. (37) in “Methods”). The $1/N$ scaling is therefore robust under individual decoherence of the components, an eventually increased prefactor not withstanding. This is corroborated by further exact results for a pure interaction with decoherence added to all $\mathcal{S}_i$ or to $\mathcal{R}$ (see “Supplementary Discussion”), and by the example of superradiance below.

Measuring the length of a cavity

As example of an application, we now show how to measure the relative change of length $\delta L/L$ of a cavity with an uncertainty of order $1/N$ with an initial product state of N quantum resources. We first consider unitary evolution.

Let N two–level atoms or ions (N even, ground and excited states $|0\rangle_i$, $|1\rangle_i$ for atom i, $i = 1, \ldots, N$) be localized in a cavity, and resonantly coupled with real coupling constants g_i to a single e.m. mode of the cavity of frequency ω and annihilation operator a (see Fig. 1), interaction hamiltonian $H_I = \sum_{i=1}^{N} g_i \left(\sigma_-^{(i)} a^\dagger + \sigma_+^{(i)} a\right)$, where $\sigma_-^{(i)} = |0\rangle_i\langle 1|_i$, $\sigma_+^{(i)} = |1\rangle_i\langle 0|_i$. Due to the spatial dependence of the e.m. mode in resonance with the atoms, the g_i depend on the position z_i of the atoms along the cavity axis and on the length L of the cavity (the waist of the mode is taken to be much larger than the size of the atomic ensemble),

$$g_i = \sqrt{\frac{\hbar\omega}{\epsilon_0 V}}\, \sin(k_z z_i)\, \boldsymbol{\epsilon}\cdot\mathbf{d}\,, \tag{15}$$

where $k_z = \pi n_z/L$, ϵ_0 denotes the dielectric constant of vacuum, $V = LA$ the mode volume (with an effective cross–section A), $\boldsymbol{\epsilon}$ the polarization vector of the mode, and $\mathbf{d}$ the vector of electric dipole transition matrix elements between the states $|0\rangle_i$ and $|1\rangle_i$, taken identical for all atoms.

If all g_i are identical, we obtain the Tavis-Cummings model [27]. Here we consider the situation where the atoms can be grouped into *two sets* with $N/2$ atoms each and coupling constants G_1 in the first set ($i \in \{1\ldots, N/2\}$), and G_2 in the second set ($i \in \{N/2+1, \ldots, N\}$). One way of obtaining two coupling constants may be to trap the atoms in two two–dimensional lattices perpendicular to the cavity axis (see Fig. 1). Note that it is not necessary to locate the atoms within a quarter wave-length of each other in order to obtain a DFS, as would be necessary without the cavity [28]. Distances which are integer multiples of the wave-length work just as well. In (15) we have neglected the transversal dependence of the mode, assuming that the atoms are localized at a distance from the cavity axis much smaller than the waist of the mode. However, this is for a computational convenience only. The initial product DFS states also exist if there is a radial variation of the g_i, but describing the dynamics would become much more complicated as it would depend on all the different g_i. Assuming two different sets of coupling constants, the system is described by eq. (1), where we identify a *pair* of atoms $(i, i+N/2)$ with subsystem $\mathcal{S}_i$, $i = 1, \ldots, N/2 \equiv N_p$, and the resonant cavity mode with the quantum bus $\mathcal{R}$. The free hamiltonian H_0 consists of the energy of all atoms, $H_i = (\omega/2)(\sigma_z^{(i)} + \sigma_z^{(i+N/2)})$, and the energy of the cavity mode, $H_R = \omega a^\dagger a$.

An expansion of the g_i about L for a small change δL allows one to write the coupling in the form of (1) with $S_{i,1}(x) = g\left((1+x)\sigma_-^{(i)} + (1-x)\sigma_-^{(i+N/2)}\right)$, $S_{i,2} = S_{i,1}^\dagger$, and $R_1 = a^\dagger$, $R_2 = a$, and $x \propto \delta L/L$ (see SI for the prefactor). For notational simplicity we restrict ourselves to the case where for $x = 0$ the couplings are the same for the two sets, $G_1 = G_2 = g$, but this is by no means necessary for the method to work.

A convenient basis for a pair of atoms is given by the “singlet” and “triplet” states $\{|s\rangle, |t_-\rangle, |t_0\rangle, |t_+\rangle\}$ with $|s\rangle = (|01\rangle - |10\rangle)/\sqrt{2}$, $|t_-\rangle = |00\rangle$, $|t_0\rangle = (|01\rangle + |10\rangle)/\sqrt{2}$, and $|t_+\rangle = |11\rangle$. As initial state of all $\mathcal{S}_i$ and $\mathcal{R}$ we take



FIG. 1: Scheme for measuring the change of the length of an optical cavity. N atoms (or ions) are trapped at fixed positions in two 2D optical lattices perpendicular to the cavity axis. A dipole transition of the atoms is in resonance with a single, leaky cavity mode. The atoms are initially prepared in a dark state in which destructive interference prevents the photons from being transferred from the atoms to the cavity mode. When the cavity length L changes by a small amount δL, the true dark states evolve, and the initial state is exposed to collective decoherence, detectable by photons leaking out through the semi-reflecting mirror at a rate proportional to N^2. This allows to measure $\delta L/L$ with a Heisenberg limited uncertainty of order $1/N$, even if the initial dark state is a product state.

the product state (7) with $N \to N/2$, and $|\varphi\rangle = (|t_-\rangle + |s\rangle)/\sqrt{2}$, and $|\xi\rangle = |0\rangle$ for a cavity mode in the vacuum state. We obtain a time-independent $K(t_1, t_2) = g^2\left(N_p + N_p^2\right)/2$, and from (6)

$$\delta x_{\min} = \frac{\sqrt{2}}{\sqrt{M} g t \sqrt{2N + N^2}}\,, \tag{16}$$

which clearly scales as $1/N$ for $N \gg 1$. One might argue that a small amount of entanglement is present in $|\varphi\rangle$, but the size of the cluster of atoms all entangled with each other (i.e. a pair of atoms) is independent of N, such that it is legitimate to consider a pair of atoms as individual subsystem, and it is a product state of these subsystems that we consider. In SI we show that the product state can be prepared by letting the atoms interact pairwise.

The initial state contains half a photon per atom. For a generic state the excitations stored in the atoms would start oscillating between the cavity mode and the atoms. However, for $x = 0$ our initial state is a "dark state", as destructive interference prevents the transfer of the photon from any pair of atoms to the cavity. When x deviates from zero, the perfect cancellation in the destructive interference is broken, and photons get transferred to the cavity.

Measuring the number of photons constitutes an optimal measurement in the sense that the bound (16) is reached. To see this, we identify $A = a^\dagger a$ in eq. (13). This leads in a straightforward manner to $\delta x = \frac{\sqrt{2}}{\sqrt{M} g t\, N}$, which agrees with (16) for $N \gg 1$, including the prefactor. After what was said in Sec. "Decoherence", it is clear that adding independent decoherence to all subsystems does not change the $1/N$ scaling of $\delta x_{\min}$. We now show this explicitly by considering the situation of very strong damping of the cavity mode, the superradiant regime.

The framework of Sec. "Decoherence" is suited for this analysis, but we adopt the well-developed theory of superradiance [29–32] to give an independent demonstration that $\delta x_{\min}$ scales as $1/N$. Decoherence arises because of two processes: Each atom can undergo spontaneous emission with rate Γ, due to its coupling to a continuum of additional e.m. modes. The damping of the cavity mode arises from the escape of photons with a rate 2κ through one of the mirrors. In the notation of eq.(14), and identification of a pair of atoms $(i, i + N/2)$ with $\mathcal{S}_i$, the generators Λ_i and Λ_R for these two processes read [29–32]

$$\begin{aligned} \Lambda_i X &= \frac{\Gamma}{2}\left([\sigma_-^{(i)} X, \sigma_+^{(i)}] + [\sigma_-^{(i+N/2)} X, \sigma_+^{(i+N/2)}] + h.c.\right)\,, && (17) \\ \Lambda_R X &= \kappa\left([aX, a^\dagger] + h.c.\right)\,. && (18) \end{aligned}$$

Superradiance occurs in the overdamped regime $\Gamma \ll g\sqrt{N} \ll \kappa$, where a photon transferred to the cavity leaves the cavity before it can feed itself back to the atoms, but induces emission in other atoms while in the cavity mode. Cavity decay is then the by far dominant process. We will therefore start by neglecting Γ, but treat spontaneous emission in SI. The population of the cavity follows the occupation of the atoms adiabatically, and one can eliminate the cavity mode. This leads to the well-known and, for $x = 0$, experimentally verified master equation of superradiance [29–34] for the reduced density matrix ρ_s of the atoms in the interaction picture,

$$\frac{d}{dt}\rho_s(t) = L_I(x)[\rho_s(t)] \equiv \gamma\left([J_-(x)\rho_s(t), J_+(x)] + [J_-(x), \rho_s(t)J_+(x)]\right). \tag{19}$$

The collective generators $J_\pm$ are $J_-(x) = \sum_{i=1}^{N/2} S_{i,1}(x) = \sum_{i=1}^{N/2}\left((1+x)\sigma_-^{(i)} + (1-x)\sigma_-^{(i+N/2)}\right)$, $J_+(x) = J_-^\dagger(x)$. The rate $\gamma = g^2/\kappa$ is independent of N. Collective decoherence is a two-stage process here, as photons stored in the atoms first need to be transferred to the cavity mode before they can leave the system. The dark states of Sec. "Measuring the length of a cavity" are therefore decoherence-free states. There is a large decoherence-free subspace (DFS) containing $\binom{N}{N/2} \sim 2^N/\sqrt{N}$ DF states, including a $2^{N/2}$ dimensional subspace $\bigotimes_{l=1}^{N/2}\{|t_-\rangle_l, |s\rangle_l\}$ in which the pair formed by the atoms l and $l+N/2$ can be in a superposition of $|t_-\rangle_l$ and $|s\rangle_l$ [24, 35]. A DFS of the same dimension also exists for non-identical couplings, but the coefficients in the linear combination of the singlet state need to be adapted accordingly, $|s\rangle_l \to \frac{1}{\sqrt{|G_1|^2+|G_2|^2}}(G_1|0\rangle_l|1\rangle_{l+N/2} - G_2|1\rangle_l|0\rangle_{l+N/2})$. If after preparing the atoms in a DFS state corresponding to the initial couplings $G_1^{(0)}$, $G_2^{(0)}$ the length L of the cavity changes slightly, the coupling constants will evolve, $G_I^{(0)} \to G_I$, $I = 1, 2$, and so will the DFS. Photons will leak out of the cavity as the original state becomes exposed to decoherence.

There is a well-known connection between the photon statistics in the cavity mode and the excitation of the atoms, derived in [30] for all couplings identical,

$$\langle a^{\dagger m}a^m(t)\rangle = 2m\left(\frac{g}{\kappa}\right)^{2m}\int_0^t ds\,\kappa\, e^{-2m\kappa s}(e^{\kappa s}-1)^{2m-1}\langle J_+^m J_-^m(t-s)\rangle. \tag{20}$$

One checks that this relation remains valid for small asymmetries $x \neq 0$. Thus, instead of the number of photons in the cavity for a given value x one can calculate the excitation of the atoms, where, however, the observable itself becomes a function of x, $J_+^m(x)J_-^m(x)$. For the initial product state (7) considered above (with $N \to N_p$, $|\varphi\rangle = (|t_-\rangle + |s\rangle)/\sqrt{2}$, and $|\xi\rangle = |0\rangle$) we have from (19),

$$\begin{aligned}
\langle J_+J_-(t)\rangle &= x^2\left(\frac{N_p}{2} + \frac{N_p^2}{2}\right) - \gamma t\left(3N_p^2 + N_p^3\right)(x^2 + x^4) + \mathcal{O}(t^2), && (21)\\
\langle J_+^2 J_-^2(t)\rangle &= \left(\frac{N_p}{2} - \frac{5N_p^2}{4} + \frac{N_p^3}{2} + \frac{N_p^4}{4}\right)x^4 \\
&+ 2\gamma t\Bigg[\left(-N_p - \frac{5}{2}N_p^2 + \frac{11}{2}N_p^3 - \frac{3}{2}N_p^4 - \frac{1}{2}N_p^5\right)x^4 \\
&+ \left(-3N_p + \frac{5}{2}N_p^2 + \frac{7}{2}N_p^3 - \frac{5}{2}N_p^4 - \frac{1}{2}N_p^5\right)x^6\Bigg] + \mathcal{O}(t^2). && (22)
\end{aligned}$$

Equation (20) is in principle valid only in the Markovian regime $t \gg 1/\kappa$, if $\langle J_+^m J_-^m(t-s)\rangle$ is obtained from the solution of the Markovian superradiance master equation (19). However, the initial behavior of $\langle a^\dagger a(t)\rangle$, $\langle a^\dagger a(t)\rangle \simeq g^2t^2\langle J_+J_-(0)\rangle$, is entirely determined by the value of $\langle J_+J_-(t)\rangle$ at $t = 0$, i.e. the question of the Markovian approximation of the dynamics of $\langle J_+J_-(t)\rangle$ does not arise, and eq. (20) can therefore be used to calculate $\langle n_{\rm ph}(x,t)\rangle$ for short times up to order t^2. From $\langle J_+J_-(0)\rangle$ one finds immediately

$$\langle n_{\rm ph}(x,t)\rangle = \langle a^\dagger a(x,t)\rangle = \frac{1}{2}g^2t^2x^2\left(N_p(N_p+1)\right) + \mathcal{O}(t^3). \tag{23}$$

At this order κ does not intervene yet, as initially the cavity mode is in the vacuum state. The quadratic initial increase of $\langle n_{\rm ph}\rangle$ reflects the beginning of a Rabi oscillation between the excited atoms and the cavity mode. We expect this result therefore to be valid as long as $\langle n_{\rm ph}(x,t)\rangle \lesssim 1$. Equation (23) agrees identically with the result one finds from the approach in Sec. "Decoherence" (see eq.(37) in "Methods"). The fluctuations of $n_{\rm ph}$ are obtained from $\langle n_{\rm ph}(x,t)^2\rangle = \langle a^{\dagger 2}a^2(x,t)\rangle + \langle n_{\rm ph}(x,t)\rangle$. Together with (20) one gets for $\langle n_{\rm ph}(x,t)\rangle \lesssim 1$, $\langle \Delta n_{\rm ph}^2(x,t)\rangle \simeq \langle n_{\rm ph}(x,t)\rangle$ with corrections of order $(g/\kappa)^4$. From eqs.(4) and (23) we find $\delta x = \frac{\sqrt{2}}{\sqrt{M}gt\sqrt{N(N+2)}} \simeq \frac{\sqrt{2}}{\sqrt{M}gtN}$, which is identical to the minimal possible uncertainty, eq.(16) for $N \gg 1$. The validity of the short time expansions (21,22) is limited to $N\gamma t \ll 1$, as can be seen from comparing the first order term with the zeroth order term. Inserting (21) in (20) gives therefore an analytical prediction of $\langle n_{\rm ph}(t)\rangle$ valid for $g/\kappa \ll gt \ll \kappa/(Ng)$, in addition to the small time result (23) for $gt \ll g/\kappa$. The agreement of $\langle n_{\rm ph}(t)\rangle$ based on (21) with the result from simulating (19) can be further improved by re-exponentiating $\langle J_+ J_-(t)\rangle$ according to $a+bt \simeq a\exp(b/at)$, before inserting it in (20). The limitation of validity of the small-time expansion does not pose a serious restriction in the bad cavity limit $\kappa \gg g$, nor does it imply that the $1/N$ scaling of the sensitivity breaks down beyond that regime. A full theoretical analysis for longer times will have to include the calculation of the superradiant propagator with broken $SU(2)$ symmetry, however. For $t \simeq 1/\kappa$ a non-Markovian description of superradiance is called for, which is beyond the scope of the present investigation.



FIG. 2: Mean photon number and uncertainty δx of the change of length of the cavity **(a)**: Mean photon number $\langle n_{\rm ph}\rangle$ as a function of dimensionless time gt (where g is the coupling constant of the atoms to the cavity mode) for $N = 2, 4, 6, 8, 10, 12$ (black, red, green, blue, brown, violet), in units of x^2 for $x = 0.1$, and with photon escape rate from cavity $\kappa = 5g$, obtained through numerical simulation of superradiance using a stochastic Schrödinger equation. The dashed lines with corresponding colors are analytical results valid up to $Ng^2t/\kappa \sim 1$ (see eqs. (20),(21)). **(b)**: Uncertainty δx (see eq. (4)) based on $A = n_{\rm ph}$ as function of N for $gt = 0.0485$ and $x = 0.01$. Numerical results (circles) show the same $1/N$ scaling as the ideal lower bound (red dashed line), eq. (16), with slightly increased prefactor. Green continuous line is an analytical prediction based on an expansion of $\langle n_{\rm ph}(t)\rangle$ for small gt.

Figure 2 shows that $\langle n_{\rm ph}(t)\rangle$ obtained numerically by simulating (19) through an equivalent stochastic Schrödinger equation (SSE), and integration of $\langle J_+ J_-(t)\rangle$ according to eq. (20), agrees well with the result based on (21) for $gt \ll g/\kappa$ and $g/\kappa \ll gt \ll \kappa/(Ng)$. The SSE for real $\psi(t)$ reads

$$d\psi(t) = D_1(\psi(t))\,dt + D_2(\psi(t))\,dW(t)\,, \tag{24}$$

with $D_1(\psi) = \gamma\left(2\langle J_-\rangle_\psi J_- - J_+J_- - \langle J_-\rangle_\psi^2\right)\psi$, and $D_2(\psi) = \sqrt{2\gamma}\,(J_- - \langle J_-\rangle_\psi)\,\psi$, where $dW(t)$ is a Wiener process with average zero and variance dt, and $\langle J_-\rangle_\psi = \langle\psi|J_-|\psi\rangle$ [26]. We used 2000 equidistant time steps in the time interval $t = 0, \ldots, 20/g$, 20 random realizations of the process for the simulation of $\langle n_{\rm ph}(t)\rangle$, and 400 realizations for the calculation of δx. Figure 2 also shows δx calculated from the numerical data for $\langle \Delta n_{\rm ph}^2(t)\rangle^{1/2}$ and $\langle n_{\rm ph}(x,t)\rangle$ through eq.(4), together with the fundamental lower bound $\delta x_{\rm min}$, eq.(16). We see that at $gt = 0.0485$, δx follows the optimal $1/N$ scaling with only slightly increased prefactor.

We emphasize that $n_{\rm ph}$ allows to measure $\delta L/L$, not just to detect a change of L. Eqs.(20) and (21) relate $\langle n_{\rm ph}\rangle$ to x, and, unless the two lattices are situated at anti-nodes of the mode, the relation between $\delta G \equiv (G_1 - G_2)/2 = gx$ and $\delta L/L$ is linear to lowest order and independent of N: If we choose the position of the atoms such that $z_2 - z_1 = m\lambda$ with $n_z - 1 \geq m \in \mathbb{N}$ we have

$$x = \frac{\delta G}{g} = m\pi \cot\left(\frac{n_z \pi z_1}{L}\right)\frac{\delta L}{L}.$$

Therefore, the measurement of $\langle n_{\rm ph}\rangle$ allows the measurement of $\delta L/L$. Several other practical questions, e.g. the preparation of the initial state, and the robustness of the method with respect to fluctuations of the coupling constants, spontaneous emission, and errors in the preparation of the initial state, are addressed in SI. The superradiant regime has the advantage of providing direct access to the number of photons in the cavity. The average number of photons outside is simply obtained by integrating $2\kappa\langle a^\dagger a(t)\rangle$ up to time t, as the photon escape rate is proportional to the average photon number inside the cavity [30]. The results for the scaling of δx with N based on $A = a^\dagger a$ are therefore unaffected by detecting the photons that leave the cavity. Measuring the number of photons amounts to monitoring the decoherence dynamics, and we have thus an example where the parametric dependence of a collective decoherence process allows to achieve the Heisenberg limit with an initial product state.

Discussion

Our results may seem to conflict with the well-known theorem [4] that for unitary evolution of N independent quantum systems in an initial product state at best a scaling $\delta x_{\min} \sim 1/\sqrt{N}$ is possible. To see that there is no contradiction, it is helpful to consider the simple case where $H'(x) \equiv dH(x)/dx$ and $H(x)$ commute, $[H(x), H'(x)] = 0$. One then easily shows that to lowest order in dx, $\rho(x,t) = \exp(-\mathrm{i}H(x)t)\rho(0)\exp(\mathrm{i}H(x)t)$ (with $\hbar = 1$) and $\rho(x,t) + d\rho$ are related by a unitary transformation with generator $\hat{h} = H'(x)\,t$. Let us furthermore restrict ourselves to a single operator per subsystem, i.e. $\nu = 1$ only, and to the linear x dependence $S_{i,1}(x) = xS$ for all i, and $R_1 = R$. A few lines of calculation lead to

$$\langle \Delta \hat{h}^2 \rangle = \left(N \langle \Delta S^2 \rangle \langle R^2 \rangle + N^2 \langle S \rangle^2 \langle \Delta R^2 \rangle \right) t^2 \,, \tag{25}$$

for an initial product state, $\rho(0) = |\psi_0\rangle\langle\psi_0|$ with $|\psi_0\rangle$ from eq.(7). All expectation values of S in (25) are in state $|\varphi\rangle$, those of R in the state $|\xi\rangle$. Inserting (25) into (3) and (2), we find that for $N \gg 1$ and $\langle S \rangle^2 \langle \Delta R^2 \rangle \neq 0$,

$$\delta x_{\min} = \frac{1}{2\sqrt{M} t |\langle S \rangle| \langle \Delta, R^2 \rangle^{1/2}} \frac{1}{N}$$

i.e. the Heisenberg limit $\delta x_{\min} \sim 1/N$ can be achieved with an initial product state. Clearly, for the case considered above the unitary transformation generated by $H_0 \equiv \sum_i H_i + H_R$ is not necessary in order to achieve the $1/N$ scaling. We therefore simplify the reasoning further by considering the case $H_0 = 0$. We are then left with a pure interaction,

$$H(x) = H_I(x) = x \sum_i S_i \otimes R \,. \tag{26}$$

But this is *not* a hamiltonian of the form $H(x) = x \sum_i h_i$ required by the theorem in [4]. In our case all subsystems couple in a non-trivial fashion to the common quantum bus $\mathcal{R}$ and are therefore not independent. This turns out to be the decisive difference. The SQL can be recovered for the standard situation of N *independent* subsystems through a R that acts only trivially on $\mathcal{R}$, i.e. $R = \mathbf{1}$, such that $\langle \Delta R^2 \rangle = 0$, and thus $\delta x_{\min} = 1/(2\sqrt{MN} t \langle \Delta S^2 \rangle^{1/2})$. This makes obvious the rather ironic fact that quantum fluctuations in $\mathcal{R}$ help and are necessary to achieve the $1/N$ scaling. The prefactor of the $1/N$ behavior is smallest for an initial state with an equal weight superposition of the eigenstates of R pertaining to its largest and lowest eigenvalues $r_{\min}$ and $r_{\max}$, in which case $\langle \Delta R^2 \rangle^{1/2} = |r_{\max} - r_{\min}|/2$.

This simple example also allows to corroborate that in order to achieve the $1/N$ scaling one need not measure the $\mathcal{S}_i$ *at all*, and almost any measurement on $\mathcal{R}$ suffices. Consider an initial product state $|\psi_0\rangle = \otimes_{i=1}^{N} |s_i\rangle_i |\xi\rangle$, with $S_i|s_i\rangle = s_i|s_i\rangle$, and $|\xi\rangle = \sum_m d_m |r_m\rangle$ for $R|r_m\rangle = r_m|r_m\rangle$. We then have

$$|\psi(t)\rangle = \sum_m d_m e^{-\mathrm{i}x \sum_i s_i r_m t} \bigotimes_{i=1}^{N} |s_i\rangle \otimes |r_m\rangle \,. \tag{27}$$

Let A be an observable on $\mathcal{R}$ which does not commute with R, i.e. there are at least two eigenstates $|r_0\rangle$ and $|r_1\rangle$ such that $\langle r_0|A|r_1\rangle \neq 0$. It is sufficient to consider an initial state of $\mathcal{R}$ which is a superposition of these two states, e.g. we may take $d_0 = d_1 = 1/\sqrt{2}$, and an observable $A = |r_0\rangle\langle r_1| + |r_1\rangle\langle r_0|$. If all subsystems $\mathcal{S}_i$ are prepared in the same state with $s_i = s$, one finds $\langle A(t) \rangle = \cos(xNs(r_0 - r_1)t)$ and $\langle \Delta A^2(t) \rangle = \sin^2(xNs(r_0 - r_1)t)$. Inserted in eq. (4) this leads to the exact result

$$\delta x = \frac{1}{N|s||r_0 - r_1|t} \,, \tag{28}$$

valid for all x. This shows that the $1/N$ scaling can be reached by measuring almost any observable of $\mathcal{R}$, as long as it does not commute with R. Furthermore, eq. (27) allows a simple quantum information theoretical explanation of the effect: The final state reflects the accumulated phase from the interaction of all the systems $\mathcal{S}_i$ with the common quantum bus $\mathcal{R}$. Figure 3 shows an equivalent quantum circuit that reproduces state (27). One subsystem $\mathcal{S}_i$ after another imprints the same phase on the components of the state of $\mathcal{R}$. Equation (27) also makes obvious that a measurement of the $\mathcal{S}_i$ alone does not allow to achieve the $1/N$ scaling, as the state of $\mathcal{R}$ will collapse on a single



FIG. 3: Quantum circuit that reproduces the interaction hamiltonian $H_I(x) = x\sum_{i=1}^N S_i \otimes R$. N quantum systems $\mathcal{S}_i$ prepared all in the eigenstate $|s\rangle$ of S_i, $S_i|s\rangle = s|s\rangle$ in eq. (26), lead to a total accumulated relative phase between states of the quantum bus $\mathcal{R}$ that is proportional to xN. This allows a measurement of x with a precision that scales as $1/N$, even though the initial state is a product state, by measuring any observable A on $\mathcal{R}$ alone that does not commute with R.

state $|r_m\rangle$, and one only gets an irrelevant global phase. Thus, measuring the quantum bus is not only sufficient, but also necessary for the $1/N$ scaling of δx. We also see that the measurement of $A = |r_0\rangle\langle r_1| + |r_1\rangle\langle r_0|$ is optimal if r_0, r_1 correspond to the smallest and largest eigenvalues of R, respectively.

In [13] an adaptive measurement technique was demonstrated that allows one to achieve Heisenberg-limited uncertainty by using only an initial product state. The method is based on phase estimation [36], but instead of using a NOON state of N photons, independent photons were passed N times through the same phase shifter. This amplifies the phase by a factor N, but it was shown that in the presence of losses the scaling of the sensitivity with N is at most improved by a constant factor [37] compared to the classical case for $N \to \infty$. A common feature of both phase estimation and our method is that a measurement is performed on a common quantum system that interacts with all other quantum systems. However, our method is more general. It incorporates decoherent and unitary evolutions in the same framework, and allows one to use collective decoherence as a signal. Secondly, phase estimation was developed for a multi-qubit system with controlled, sequentially turned on interactions, and an x-dependence in the free evolution. Hamiltonian (1) on the other hand can be used to describe substantially more complex systems, with possibly non-trivial dynamics in the absence of the collective interaction, and with interactions that do not commute with the Hamiltonians of the free constituents. Furthermore, the interaction is simultaneous such there is no bandwidth penalty in the accumulation of the phase, nor is there a need to re-sample a phase shift many times.

Our method requires a collective interaction between N separable quantum systems and a common quantum bus, an initial noise-less state in the sense discussed above, and the possibility to measure at least part of the quantum bus. In the case of incomplete measurement of the quantum bus this implies the need of a collective decoherence process with a decoherence-free initial state. Besides atoms in a cavity one might consider circuit-QED systems [38], trapped ions coupled to a common phonon mode [39], or quantum dots coupled to micro-resonators [40] or to photonic crystals [41]. Both unitary evolution or a decoherence process can be useful, as long as the collective interaction between the N quantum resources and a common quantum bus depends on the parameter x to be measured.

To summarise, we have developed a general theory of collectively enhanced quantum measurements based on the interaction of N quantum systems with a common "quantum bus". The latter can be a simple quantum system, or an environment with many degrees of freedom to which we have only partial access. We have shown that if the collective interactions depend on a parameter x, the Heisenberg limit (i.e. a $1/N$ scaling of the uncertainty of x) can be reached with an initial product state, and by measuring almost any observable of the quantum bus. We have used quantum parameter estimation theory to establish that a $1/N$ scaling of the uncertainty is indeed optimal in this setup. We have given a simple quantum-information theoretical interpretation of the effect, and we have analysed in detail a possible experimental implementation of the measurement of the change of the length of a cavity with an uncertainty that scales as $1/N$.

The proposed measurement principle offers an attractive way out of the dilemma of ubiquitous decoherence that has so far plagued quantum enhanced measurements: First of all, there is no need to build highly entangled states which are extremely fragile under decoherence for large N. Simple product states will do, and decoherence of some parts of the system does not affect the $1/N$ scaling of the minimal uncertainty. Secondly, parameter-dependent collective decoherence is covered itself by our new measurement principle. Indeed, decoherence is a process in which quantum interference effects can play an important role. This is exemplified by the very existence of DFS, and can lead to exquisite sensitivity when a DFS is disturbed. Instead of trying to suppress decoherence at all costs, one might therefore be better off exploiting its parametric dependence.

Methods

Bures distance for unitary evolution

The state vector $|\psi_I(x,t)\rangle$ in the interaction picture obeys the time-dependent Schrödinger equation

$$\mathrm{i}\frac{\partial}{\partial t}|\psi_I(x,t)\rangle = H_I(x,t)|\psi_I(x,t)\rangle\,, \tag{29}$$

with the interaction hamiltonian

$$\begin{aligned} H_I(x,t) &\equiv \sum_{i,\nu} S_{i,\nu}(x,t)\otimes R_\nu(t)\,, &(30)\\ S_{i,\nu}(x,t) &= e^{\mathrm{i}H_i t}S_{i,\nu}(x)e^{-\mathrm{i}H_i t},\quad R_\nu(t) = e^{\mathrm{i}H_R t}R_\nu e^{-\mathrm{i}H_R t}\,. &(31)\end{aligned}$$

The general solution of (29) is given by $|\psi_I(x,t)\rangle = \mathrm{T}\exp\left[-\mathrm{i}\int_0^t H_I(x,t')dt'\right]|\psi_0\rangle$, where T denotes the time-ordering operator. To second order in the perturbation $H_I(x,t)$, the overlap between $|\psi_I(x,t)\rangle$ and $|\psi_I(x+dx,t)\rangle$ reads

$$\begin{aligned}\langle\psi_I(x+dx,t)|\psi_I(x,t)\rangle &= 1 + dx\Big[\mathrm{i}\int_0^t\langle H_I'(x,t_1)\rangle dt_1 + \int_0^t\int_0^{t_1}\langle[H_I'(x,t_1),H_I(x,t_2)]\rangle dt_1dt_2\Big]\\ &+\frac{1}{2}dx^2\Big(\mathrm{i}\int_0^t\langle H_I^{''}(x,t_1)\rangle dt_1 + \int_0^t\int_0^{t_1}\langle[H_I^{''}(x,t_1),H_I(x,t_2)]\rangle dt_1dt_2\\ &-2\int_0^t\int_0^{t_1}\langle H_I'(x,t_1)H_I'(x,t_2)\rangle dt_1dt_2\Big)\,, \qquad (32)\end{aligned}$$

with all expectation values with respect to $|\psi_0\rangle$. We assume that the derivatives of $H_I(x)$ with respect to x are hermitian operators, in which case the term linear in dx is purely imaginary. The lowest order term in the squared overlap is then of order dx^2. One finds in a straightforward manner the squared Bures distance (5).

Decoherence of subsystems

Markovian decoherence of the $\mathcal{S}_i$ and $\mathcal{R}$ on top of the unitary evolution generated by $H(x)$ can be described by eq. (14). The single system dynamics (i.e. all $\mathcal{S}_i$ and $\mathcal{R}$ taken separately, $L_I=0$), can be solved formally by exponentiating the Liouvillians. We will again treat $L_I(x)$ in perturbation theory. The density matrix $W_I(t)$ of $\mathcal{S}$ and $\mathcal{R}$ in the interaction picture is related to the one in the Schrödinger picture, $W(t)$, by

$$W(t) = e^{-\mathrm{i}(L_0+\mathrm{i}\Lambda_s+\mathrm{i}\Lambda_r)t}W_I(t)\equiv e^{-\mathrm{i}Kt}W_I(t)\,, \tag{33}$$

and obeys the master equation $\dot{W}_I(t) = -\mathrm{i}L_I(x,t)W_I(t)$. With eq. (33) we have defined the free propagator $P_F(t)\equiv e^{-\mathrm{i}Kt} = \otimes_i e^{-\mathrm{i}K_i t}\otimes e^{-\mathrm{i}K_R t}\equiv\otimes_i P_i(t)\otimes P_R(t)$, and $L_I(x,t) = P_F(-t)L_I(x)P_F(t)$ is the interaction Liouvillian in the interaction picture. We decompose furthermore $L_I = \sum_{k=1}^N L_{I,k}$, with $L_{I,k}X = [H_{I,k},X]$. To second order in L_I we have

$$\begin{aligned}\langle A(t)\rangle &= \mathrm{tr}\Big\{A\Big(P_F(t) - \mathrm{i}\sum_k\int_0^t P_F(t-t_1)L_{I,k}(x)P_F(t_1)\,dt_1 &(34)\\ &-\sum_{k,k'}\int_0^t dt_1\int_0^{t_1}dt_2 P_F(t-t_1)L_{I,k}(x)P_F(t_1-t_2)L_{I,k'}(x)P_F(t_2) + \mathcal{O}(L_{I,k}^3)\Big)W(0)\Big\}\,.\end{aligned}$$

With an initial product state, $W(0) = \otimes_{i=1}^N\rho_i(0)\otimes\rho_r(0)$, we obtain the zeroth order term $\langle A(t)\rangle_0\equiv\mathrm{tr}(AP_F(t)W(0)) = \mathrm{tr}(\otimes_i P_i(t)[\rho_i(0)]\otimes AP_R(t)[\rho_R(0)]) = \mathrm{tr}_R(A\rho_R(t))$, as all propagators are trace-preserving, and $\rho_R(t)\equiv P_R(t)[\rho_R(0)]$. Similarly, by explicitly writing $L_{I,k}X = \sum_\nu[S_{k,\nu}R_\nu,X]$, we obtain the first order term

$$\begin{aligned}\langle A(t)\rangle_1 &\equiv -\mathrm{i}\sum_k\mathrm{tr}\left(A\int_0^t P_F(t-t_1)L_{I,k}P_F(t_1)\,dt_1W(0)\right) &(35)\\ &= -\mathrm{i}\sum_{k,\nu}\int_0^t dt_1\langle S_{k,\nu}(t_1)\rangle\mathrm{tr}_R\left(AP_R(t-t_1)[R_\nu,\rho_R(t_1)]\right)\,, &(36)\end{aligned}$$

where $\langle S_{k,\nu}(t)\rangle = \mathrm{tr}_k S_{k,\nu} P_k(t)[\rho_k(0)]$. We generalise the second simplifying assumption in Sec. "Measuring the quantum bus" to $P_R(t)[\rho_R(0)] = \rho_R(0)$, and $\mathrm{tr}_R(AP_R(t)[X]) = f_A(t)\mathrm{tr}_R(AX)$ with some function $f_A(t)$ [30]. This implies that the initial state is decoherence-free concerning the decoherence of $\mathcal{R}$ alone. This is a natural assumption for the state of an environment initially in thermal equilibrium, or for a quantum bus in its ground state, such as an initially empty cavity mode (see the example of superradiance). We then have again $\langle A(t)\rangle_1 = 0$. To second order in the interaction we find

$$\begin{aligned}
\langle A(t)\rangle &= \langle A(0)\rangle_0 \\
&- \sum_{\nu,\mu}\int_0^t dt_1 \int_0^{t_1} dt_2 \Big\{ N\Big(C^{(1)}_{S\mu\nu}(t_1,t_2) C^{(1)}_{AR\mu\nu}(t,t_1,t_2) - C^{(2)}_{S\mu\nu}(t_1,t_2) C^{(2)}_{AR\mu\nu}(t,t_1,t_2)\Big) \\
&+ N^2 \langle S_\mu(t_1)\rangle\langle S_\nu(t_2)\rangle C^{(3)}_{AR\mu\nu}(t,t_1,t_2)\Big\}\,, \qquad (37)\\
C^{(1)}_{AR\mu\nu}(t,t_1,t_2) &= \mathrm{tr}_R\left(AP_R(t-t_1)[R_\mu, P_R(t_1-t_2)[R_\nu \rho_R(t_2)]]\right) \qquad (38)\\
C^{(2)}_{AR\mu\nu}(t,t_1,t_2) &= \mathrm{tr}_R\left(AP_R(t-t_1)[R_\mu, P_R(t_1-t_2)[\rho_R(t_2) R_\nu]]\right) \qquad (39)\\
C^{(3)}_{AR\mu\nu}(t,t_1,t_2) &= C^{(1)}_{AR\mu\nu}(t,t_1,t_2) - C^{(2)}_{AR\mu\nu}(t,t_1,t_2) \qquad (40)\\
C^{(1)}_{S\nu\mu}(t_1,t_2) &= \mathrm{tr}_k\left(S_\mu P_k(t_1-t_2)[S_\nu \rho_k(t_2)]\right) - \langle S_\mu(t_1)\rangle\langle S_\nu(t_2)\rangle \qquad (41)\\
C^{(2)}_{S\nu\mu}(t_1,t_2) &= \mathrm{tr}_k\left(S_\mu P_k(t_1-t_2)[\rho_k(t_2) S_\nu]\right) - \langle S_\mu(t_1)\rangle\langle S_\nu(t_2)\rangle\,, \qquad (42)
\end{aligned}$$

where $\rho_R(t) = P_R(t)[\rho_R(0)]$. The index k is arbitrary, $k = 1,\ldots,N$, as we have assumed all systems $\mathcal{S}_k$ identical and identically prepared. All x dependence is in the operators S_ν. Equation (37) also gives $\langle A^2(t)\rangle$ by replacing $A \to A^2$, and $\langle A(t)\rangle^2$ to order $\mathcal{O}(H_I^2)$. The equation obtained by inserting these expressions into eq. (4) generalizes the result (12) to decoherence on top of the unitary evolution considered in Sec. "Measuring the quantum bus". We see that the basic structure of the result for δx, and in particular its scaling with N is unchanged, but the expectation values and correlation functions are replaced by more complicated expressions involving in general mixed states and non-unitary evolution of individual subsystems.

Quantum parameter estimation for a Markovian master equation

In standard descriptions of decoherence one traces out the heat bath and gets a master equation for the reduced density matrix ρ_s of $\mathcal{S}$ alone. Using quantum parameter estimation theory generalized to non-unitary evolution we now show that for Markovian decoherence with an initially decoherence-free state, measuring an arbitrary x-independent observable on $\mathcal{S}$ alone gives at best a $\delta x_{\min} \sim 1/\sqrt{N}$. This corroborates the result found for unitary evolution that the important quantum system to measure is the common quantum bus $\mathcal{R}$, rather than $\mathcal{S}$.

The Markovian master equation for $\rho_s(t)$ obtained by tracing out $\mathcal{R}$ has the Lindblad-Kossakowski form

$$\dot{\rho_s}(t) = \Lambda(x)[\rho_s(t)] \equiv \gamma \sum_{\alpha=1}^{d} \left([F_\alpha(x), \rho_s(t) F_\alpha^\dagger(x)] + h.c.\right)\,, \qquad (43)$$

where we work in the interaction picture and assume that there is no additional unitary evolution. The $F_\alpha(x)$ are arbitrary linear (not necessarily hermitian) operators which have inherited the x-dependence from the interaction hamiltonian $H_I(x)$, and d is the total number of generators. Note that we can restrict ourselves to an initially pure state, as for any linear propagation one cannot do better with a mixed state than with the pure states from which it is mixed [42]. We expand the Markovian time evolution to first order in t, $\rho = |\psi\rangle\langle\psi| + t\,\Lambda(x)|\psi\rangle\langle\psi| + \mathcal{O}(t^2)$, and linearise $F_\alpha(x)$ about the value of x where we want to measure. We set that value, without restriction of generality, to zero, i.e. $F_\alpha(x) = xF'_\alpha + F_\alpha(0)$, and assume that the initial state is decoherence-free at $x = 0$. The Bures distance can still be evaluated in a straight-forward fashion as the state at $x = 0$ remains pure. One finds $ds^2\mid_{x=0} = 8\gamma t \sum_\alpha K_{|\psi\rangle}(F_\alpha^\dagger, F_\alpha)\, dx^2$. As a consequence, the ultimate quantum limit of the sensitivity with which the parameter x can be estimated from the parametric dependence of the master equation, starting from a pure state $|\psi\rangle$, reads

$$\delta x_{\min} = \frac{1}{2\sqrt{2M\gamma t}\left(\sum_{\alpha=1}^{d} K_{|\psi\rangle}(F_\alpha^\dagger, F_\alpha)\right)^{1/2}}\,. \qquad (44)$$

With $F_\alpha = \sum_{r=1}^N F_{\alpha,r}$ we obtain $K_{|\psi\rangle}(F_\alpha^\dagger, F_\alpha) = \sum_{r,s=1}^N K_{|\psi\rangle}(F_{\alpha,r}^\dagger, F_{\alpha,s})$. For an initially entangled state, $K_{|\psi\rangle}(F_\alpha^\dagger, F_\alpha)$ can be of order N^2. This can be seen from the example of the GHZ state $|\psi\rangle = (|0\dots0\rangle + |1\dots1\rangle)/\sqrt{2}$, and a single generator $F_{1,i} = \sigma_z^{(i)}$ with $\sigma_z^{(i)}$ the Pauli z matrix for subsystem i. Then $K_{|\psi\rangle}(F_1^\dagger, F_1) = N^2$, and one obtains a $1/N$ scaling of $\delta x_{\min}$, just as in the case of unitary evolution. However, if the initial state factorizes, $|\psi\rangle = \otimes_{l=1}^N |\varphi_l\rangle$, there are no correlations between different subsystems r and s, and we thus have only the sum of correlations in all subsystems, $K_{|\psi\rangle}(F_\alpha^\dagger, F_\alpha) = \sum_{s=1}^N K_{|\varphi_s\rangle}(F_{\alpha,s}^\dagger, F_{\alpha,s})$, which is at most of order N, and $\delta x_{\min}$ scales as $1/\sqrt{N}$, again just as in the case of unitary evolution. This shows once more that a measurement of an x-independent observable on $\mathcal{S}$ does not allow to do better than in the standard situation of unitary evolution of $\mathcal{S}$ without coupling to a common quantum bus.

Interestingly, superradiance *is* described by a master equation of $\mathcal{S}$ alone after tracing out the cavity mode. But a measurement on $\mathcal{R}$ (the number of photons in the cavity) translates in that case to a measurement on the $\mathcal{S}_i$ that depends itself on x. In this way it is still possible to achieve a $1/N$ scaling of δx.

The authors declare to have no competing financial interests.

Author contributions: DB conceived the original idea, worked out the general formalisms and examples, found the underlying quantum-information theoretical principle, and wrote the manuscript. JM and DB both calculated the specific example of superradiance and discussed all aspects of the work at all stages.

---

[1] Cramér, H., *Mathematical Methods of Statistics* (Princeton University Press, Princeton, NJ, 1946).

[2] Braunstein, S. L. & Caves, C. M., Statistical distance and the geometry of quantum states, *Phys. Rev. Lett.* **72**, 3439–3443 (1994).

[3] Giovannetti, V., Loyd, S., & Maccone, L., Quantum-Enhanced Measurements: Beating the Standard Quantum Limit, *Science* **306**, 1330–1336 (2004).

[4] Giovannetti, V., Lloyd, S., & Maccone, L., Quantum Metrology, *Phys. Rev. Lett.* **96**, 010401 (2006).

[5] Budker, D. & Romalis, M., Optical magnetometry, *Nature Physics* **3**, 227–234 (2007).

[6] Goda, K. *et al.*, A quantum-enhanced prototype gravitational-wave detector, *Nature Physics* **4**, 472–476 (2008).

[7] Caves, C. M., Quantum-mechanical noise in an interferometer, *Phys. Rev. D* **23**, 1693–1708 (1981).

[8] Sanders, B. C., Quantum dynamics of the nonlinear rotator and the effects of continual spin measurement, *Phys. Rev. A* **40**, 2417–2427 (1989).

[9] Boto, A. N. *et al.*, Quantum Interferometric Optical Lithography: Exploiting Entanglement to Beat the Diffraction Limit, *Phys. Rev. Lett.* **85**, 2733–2736 (2000).

[10] Mitchell, M. W., Lundeen, J. S., & Steinberg, A. M., Super-resolving phase measurements with a multiphoton entangled state, *Nature* **429**, 161–164 (2004).

[11] Leibfried, D. *et al.*, Creation of a six-atom 'Schrodinger cat' state, *Nature* **438**, 639–642 (2005).

[12] Nagata, T., Okamoto, R., O'Brien, J. L., & Takeuchi, K. S. S., Beating the Standard Quantum Limit with Four-Entangled Photons, *Science* **316**, 726–729 (2007).

[13] Higgins, B. L., Berry, D. W., Bartlett, S. D., Wiseman, H. M., & Pryde, G. J., Entanglement-free Heisenberg-limited phase estimation, *Nature* **450**, 393–396 (2007).

[14] Zurek, W. H., Decoherence and the Transition from Quantum to Classical, *Phys. Today* **44**, (10), 36–44 (1991).

[15] Giulini, D. *et al.*, *Decoherence and the Appearance of a Classical World in Quantum Theory* (Springer, Berlin, Heidelberg, 1996).

[16] Braun, D., Haake, F., & Strunz, W., Universality of Decoherence, *Phys. Rev. Lett.* **86**, 2913–2917 (2001).

[17] Strunz, W. T., Haake, F., & Braun, D., Universality of decoherence for macroscopic quantum superpositions, *Phys. Rev. A* **67**, 022101 (2003).

[18] Brune, M. *et al.*, Observing the Progressive Decoherence of the "Meter" in a Quantum Measurement, *Phys. Rev. Lett.* **77**, 4887–4890 (1996).

[19] Guerlin, C. *et al.*, Progressive field-state collapse and quantum non-demolition photon counting, *Nature* **448**, 889 – 893 (2007).

[20] Zanardi, P. & Rasetti, M., Noiseless Quantum Codes, *Phys. Rev. Lett.* **79**, 3306–3309 (1997).

[21] Braun, D., Braun, P. A., & Haake, F., Slow Decoherence of Superpositions of Macroscopically Distinct States, *Proceedings of the 1998 Bielefeld Conference on "Decoherence: Theoretical, Experimental, and Conceptual Problems", Lect. Notes Phys.* **538**, 55–66 (2000).

[22] Lidar, D. A., Chuang, I. L., & Whaley, K. B., Decoherence-Free Subspaces for Quantum Computation, *Phys. Rev. Lett.* **81**, 2594–2597 (1998).

[23] Duan, L. M. & Guo, G. C., Prevention of dissipation with two particles, *Phys. Rev. A* **57**, 2399–2402 (1998).

[24] Braun, D., *Dissipative Quantum Chaos and Decoherence*, vol. 172 of *Springer Tracts in Modern Physics* (Springer, 2001).

[25] Bengtsson, I. & Życzkowski, K., *Geometry of quantum states: an introduction to quantum entanglement* (Cambride University Press, 2006).

[26] Breuer, H.-P. & Petruccione, F., *The Theory of Open Quantum Systems* (Oxford University Press, 2006).

[27] Tavis, M. & Cummings, F. W., Exact Solution for an N-Molecule—Radiation-Field Hamiltonian, *Phys. Rev.* **170**, 379–384 (1968).

[28] Karasik, R. I., Marzlin, K.-P., Sanders, B. C., & Whaley, K. B., Multiparticle decoherence-free subspaces in extended systems, *Phys. Rev. A* **76**, 012331 (2007).

[29] Agarwal, G. S., Master-Equation Approach to Spontaneous Emission, *Phys. Rev. A* **2**, 2038 (1970).

[30] Bonifacio, R., Schwendiman, P., & Haake, F., Quantum Statistical Theory of Superradiance I, *Phys. Rev. A* **4**, 302–313 (1971).

[31] Glauber, R. J. & Haake, F., Superradiant pulses and directed angular momentum states, *Phys. Rev. A* **13**, 357–366 (1976).

[32] Gross, M. & Haroche, S., Superradiance: An Essay on the Theory of Collective Spontaneous Emission, *Phys. Rep.* **93**, 301–396 (1982).

[33] Gross, M., Fabre, C., Pillet, P., & Haroche, S., Observation of Near-Infrared Dicke Superradiance on Cascading Transitions in Atomic Sodium, *Phys. Rev. Lett.* **36**, 1035–1038 (1976).

[34] Skribanowitz, N., Herman, I. P., MacGillivray, J. C., & Feld, M. S., Observation of Dicke Superradiance in Optically Pumped HF Gas, *Phys. Rev. Lett.* **30**, 309 (1973).

[35] Beige, A., Braun, D., & Knight, P. L., Driving atoms into decoherence-free states, *New Journal of Physics* **2**, 22.1–22.15 (2000).

[36] Kitaev, A. Y., Quantum measurements and the Abelian stabilizer problem, *Electr. Coll. Comput. Complex.* **3** (1996).

[37] Kołodyński, J. & Demkowicz-Dobrzański, R., Phase estimation without a priori phase knowledge in the presence of loss, *Phys. Rev. A* **82**, 053804 (2010).

[38] Fink, J. M. *et al.*, Dressed Collective Qubit States and the Tavis-Cummings Model in Circuit QED, *Phys. Rev. Lett.* **103**, 083601 (2009).

[39] Häffner, H., Roos, C. F., & Blatt, R., Quantum computing with trapped ions, *Physics Reports* **469** (2008).

[40] Reithmaier, J. P. *et al.*, Strong coupling in a single quantum dot-semiconductor microcavity system, *Nature* **432**, 197–200 (2004).

[41] Ganesh, N. *et al.*, Enhanced fluorescence emission from quantum dots on a photonic crystal surface, *Nature Nanotechnology* **2**, 515 (2007).

[42] Braun, D., Parameter estimation with mixed quantum states, *Eur. Phys. J. D* **59**, 521–523 (2010).

Acknowledgments: We thank Peter Braun, David Guery-Odelin, Jacob Taylor, and Eite Tiesinga for useful discussions. This work was supported by the Agence National de la Recherche (ANR), project INFOSYSQQ. Numerical calculations were partly performed at CALMIP, Toulouse. J.M. thanks the Belgian F.R.S.-FNRS for financial support, and DB thanks the Joint Quantum Institute, where part of this work was done, for hospitality.

Correspondence and requests for additional material should be addressed to DB.

Supplementary Discussion for the manuscript “Heisenberg-limited sensitivity with decoherence-enhanced measurements”

Pure interaction and decoherence

We reconsider here the simple interaction-dominated model (26), but with additional decoherence of the subsystems. To simplify things and make connection to the language of quantum information theory, we consider directly the case where both the $\mathcal{S}_i$ and $\mathcal{R}$ are qubits. In the eigenbasis of S_i and R, the interaction $H_I(x)$ can be written as $H_I(x) = x\sum_{i=1}^N \sigma_z^{(i)} \otimes \sigma_z^{(0)}$, where we use the label 0 for $\mathcal{R}$. As observable on $\mathcal{R}$ we chose $A = \sigma_x$. Without decoherence one obtains the uncertainty $\delta x = 1/(2\sqrt{M}Nt)$ in agreement with (28). We will restrict ourselves to phase-flip channels acting independently on all qubits with a rate Γ,

$$\Lambda_{\sigma_z^{(i)}} X \;\equiv\; \frac{\Gamma}{2}\left([\sigma_z^{(i)} X, \sigma_z^{(i)}] + [\sigma_z^{(i)}, X\sigma_z^{(i)}]\right) . \tag{S1}$$

As before we decompose the Liouvillian $L_I(x)X = [H_I(x), X] = \sum_{i=1}^N L_{I,i}X$. One easily verifies $[\Lambda_{\sigma_z^{(i)}}, L_{I,k}] = 0$ for all i, k.

Consider first phase flips of the qubits $\mathcal{S}_i$. The equation of motion for the full density matrix $W(t)$, $\dot{W}(t) = \sum_{i=1}^N (L_{I,i} + \Lambda_{\sigma_z^{(i)}})W$, is easily solved in the computational basis, ($\mathbf{a} = a_N a_{N-1} \dots a_0$, $a_i = \pm 1$, $i = 0, \dots, N$), $W_{\mathbf{ab}}(t) = W_{\mathbf{ab}}(0) \exp\left(-\mathrm{i}\sum_{i=1}^N \left(x\left(a_i a_0 - b_i b_0\right) + \mathrm{i}\Gamma(a_i b_i - 1)\right)t\right)$. The reduced density matrix for $\mathcal{R}$ alone is therefore

$$\rho_{a_0 b_0} = \sum_{a_1,\dots,a_N=\pm 1} W_{a_N\dots a_1 a_0 a_N \dots a_1 b_0}(0) \exp\left(-\mathrm{i}\sum_{i=1}^N x a_i (a_0 - b_0) t\right) . \tag{S2}$$

We see that Γ drops out completely, and dephasing of the $\mathcal{S}_i$ has therefore no impact at all on the total accumulated phase of $\mathcal{R}$, and therefore on δx.

If phase flips occur in $\mathcal{R}$, $\dot{W} = \left(\left(\sum_{i=1}^N L_{I,i}\right) + \Lambda_{\sigma_z^{(0)}}\right)W$, we have $W_{\mathbf{ab}}(t) = W_{\mathbf{ab}}(0)\exp\left(-\mathrm{i}\left(\sum_{i=1}^N x\left(a_i a_0 - b_i b_0\right) + \mathrm{i}\Gamma(a_0 b_0 - 1)\right)t\right)$. For the initial state $|\psi_0\rangle = |+\dots+\rangle(|+\rangle + |-\rangle)/\sqrt{2}$, we find the reduced density matrix element $\rho_{+-}(t) = \exp(-2\Gamma t)\cos(2Nxt)$. Thus, the decay of $\langle\sigma_x(t)\rangle$ is independent of N, and only the prefactor in the scaling of δx with N changes.

Superradiance: practical issues

We now address several additional technical questions that are important for a possible experimental implementation of our method of measuring the length of a cavity using decoherence-enhanced measurements.

Preparation of initial state

In order to prepare the product state $|\psi_0\rangle = \left(\otimes_{i=1}^N \frac{1}{\sqrt{2}}(|t_-\rangle_i + |s\rangle_i)\right)|0\rangle$ it is helpful to use three–level atoms with a lambda structure. Let $|0\rangle$ and the additional state $|2\rangle$ be hyperfine (HF) states, and assume that their energies are sufficiently split such that only the transition $|0\rangle \leftrightarrow |1\rangle$ resonates with the cavity mode. We assume further that the second optical lattice can be moved along the cavity axis, such that controlled pairwise collisions of corresponding atoms in the two lattices can be induced. Entangled pairs of atoms in their HF split ground states can thus be created (for atoms in the same lattice this has been demonstrated experimentally, see Supplementary Reference [43] for a review). After the creation of an entangled HF state $|\psi_0'\rangle$, that differs from $|\psi_0\rangle$ by the replacement of states $|1\rangle$ by states $|2\rangle$, the second lattice is moved back to its original position. Now one can selectively excite the $|2\rangle$ states by a laser pulse in resonance with the $|2\rangle \leftrightarrow |1\rangle$ transition, that replaces the singlets in the (very long lived) HF states by the desired singlets of the $|0\rangle$ and $|1\rangle$ states and thus produce $|\psi_0\rangle$. However, as such, the method is not of much practical use yet, as it will be virtually impossible to park the second lattice at the exact position corresponding to coupling constants which render $|\psi_0\rangle$ decoherence free. The extreme sensitivity of the collective decoherence with respect to changes of the coupling constants plays against us here, and will lead to leaking of light from the cavity after the excitation $|\psi_0'\rangle \to |\psi_0\rangle$, if the exact position corresponding to $|\psi_0\rangle \in$ DFS is not achieved. But it is possible to position the second lattice at the required position with a precision of $\mathcal{O}(1/N)$ using a feed-back mechanism and a part of the quantum resources. With the atoms in the state $|\psi_0'\rangle$, do the following repeatedly in order to find the optimal position: Excite a part of the entangled HF pairs containing $\mathcal{O}(N/\ln N)$ atoms with the laser, measure $\langle n_{\mathrm{ph}}\rangle$, and use the measurement results to bracket the minimum of $\langle n_{\mathrm{ph}}\rangle$ as function of the lattice position. The minimum of $\langle n_{\mathrm{ph}}\rangle$ indicates that the position corresponding to the DFS is achieved. Using golden section search, the minimum can be bracketed to precision $1/N$ in $\mathcal{O}(\ln N)$ moves, as at each step the sensitivity of the measurement of the position of the lattice is of order $\mathcal{O}(\ln N/N) \sim \mathcal{O}(1/N)$. Once the minimum is found, excite the remaining unused pairs (there should be still a number of pairs of $\mathcal{O}(N)$) to the desired state $|\psi_0\rangle$. That state is now decoherence-free, and the system ready to detect small changes of the position of one of the mirrors. Note that for this method it is not necessary to know which exact state is produced in the controlled collisions and subsequent laser excitation.

Imperfections

Here we discuss how the most important additional noise sources affect the photon statistics. In view of eqs. (20,21), we restrict ourselves to analyzing $\alpha = -\langle \dot{J}_z(x,0)\rangle \equiv \mathrm{tr}(J_z L_I(x)[\rho_s(0)]) = 2\gamma\langle J_+J_-(x,0)\rangle$, where $J_z \equiv \frac{1}{2}\sum_{i=1}^N \sigma_z^{(i)}$ is the total population inversion of the atoms. Energy conservation shows that α can be interpreted as the initial photon escape rate from the cavity.

Spontaneous emission

It is easily verified that the spontaneous emission term in eq. (17) leads to a contribution to $\langle \dot{J}_z(0)\rangle$ that scales as $\mathcal{O}(N)$, to be compared to the term of $\mathcal{O}(N^2)$ from collective emissions, eq. (21). The modification of the initial state due to spontaneous emission is not an issue, as we consider an initial product state. Atoms which decay become simply unavailable for collective decoherence, but since their number is proportional to N this does not change the scaling of δx with N. Also, note that spontaneous emission sends photons into the entire open space but *not* into the cavity, whereas the collective emission escapes *exclusively* through the leaky cavity mirror. Therefore, in addition, the two contributions can be well separated experimentally by observing only the photons which escape through the cavity mirror.

Fluctuating coupling constants

Another obvious concern are fluctuations of the coupling constants. In order to reduce the noise of the measurement of $\langle n_{\mathrm{ph}}\rangle$, one may want to repeat the experiment M times with $M \gg 1$. The exact coupling constants might fluctuate during the averaging, e.g. due to fluctuating traps caused by vibrations in the set up. But even for perfectly stable traps, thermal motion, or even quantum fluctuations in the traps will lead to fluctuating g_i. We now show that the cost in sensitivity of these fluctuations depends on their *correlations*. Fluctuations correlated between pairs of atoms l and $l+N/2$ come at no cost, completely uncorrelated fluctuations lead back to a $\delta x \propto 1/\sqrt{N}$, and fluctuations perfectly correlated within the same lattice but not between the two lattices lead to noise indistinguishable from the signal.

To see this, consider fluctuations δg_i of the g_i about their mean values G_I, $g_i = G_1 + \delta g_i$ for $i = 1,\ldots,N/2$, $g_i = G_2 + \delta g_i$ for $i = N/2+1,\ldots,N$. The generator J_- reads then $J_- = \frac{1}{g}\sum_{i=1}^N g_i\sigma_-^{(i)}$. We introduce the correlation matrix $C_{ij} = \overline{\delta g_i \delta g_j}/g^2$, where the over-line denotes an average over the ensemble describing the fluctuations, and assume $\overline{\delta g_i} = 0$ for simplicity. Then $J_-|\psi_0\rangle = -\frac{1}{2g}\sum_{i=1}^{N/2}(g_i - g_{i+\frac{N}{2}})|t_-\rangle_i \bigotimes_{l\neq i}^{N/2}|\varphi_l\rangle_l$ and

$$\alpha \;=\; \frac{\gamma}{2g^2}\Big(\sum_{i=1}^{N/2}(g_i - g_{i+N/2})^2 + \sum_{\substack{i,j=1\\ i\neq j}}^{N/2}\frac{1}{2}\left(g_i - g_{i+N/2}\right)\left(g_j - g_{j+N/2}\right)\Big)\,. \tag{S3}$$

Equation (S3) predicts a background α_{bg} in the photon escape rate on top of the value for $\delta g_i = 0$,

$$\alpha_0 = \frac{\gamma}{4}x^2(2N + N^2)\,, \tag{S4}$$

with $x = (G_1 - G_2)/(2g)$, see eq. (21), and fluctuations $\delta\alpha_f$, $\alpha = \alpha_0 + \alpha_{\rm bg} + \delta\alpha_f$, where the average background is given by

$$\overline{\alpha}_{\rm bg} \;=\; \frac{\gamma}{2}\Big\{\sum_{i=1}^{N/2}\mathcal{C}_{ii}+\frac{1}{2}\sum_{i\neq j}^{N/2}\mathcal{C}_{ij}\Big\}\,,$$

where $\mathcal{C}_{ij} \equiv \Big(C_{ij} + C_{i+\frac{N}{2}\,j+\frac{N}{2}} - C_{i\,j+\frac{N}{2}} - C_{i+\frac{N}{2}\,j}\Big)$. The average background $\overline{\alpha}_{\rm bg}$ can be determined independently at $x = 0$, and subtracted from the signal; it does therefore not influence the sensitivity of the measurement. The remaining noise $\delta\alpha_f$ fluctuates about zero,

$$\delta\alpha_f \;=\; x\gamma\left(\frac{N}{2}+1\right)\sum_{i=1}^{N/2}\frac{\Delta g_i}{g}\,, \tag{S5}$$

where $\Delta g_i \equiv \delta g_i - \delta g_{i+N/2}$. Its standard deviation $(\overline{\delta\alpha_f^2})^{1/2} = \gamma|x|D$, where $D = (\frac{N}{2}+1)\left(\sum_{i,j=1}^{N/2}\mathcal{C}_{ij}\right)^{1/2}$ translates into additional noise in the number of detected photons. Several interesting cases can be considered:

1. Fully uncorrelated fluctuations, $C_{ij} = C_i\delta_{ij}$, where δ_{ij} stands for the Kronecker-delta: Here we get $D = (\frac{N}{2}+1)\left(\sum_{i=1}^{N/2}(C_i + C_{i+N/2})\right)^{1/2}$, which is in general of order $N^{3/2}$, and leads back to the standard quantum limit, $\delta x \sim 1/\sqrt{N}$ for large N.

2. Pairwise identical fluctuations between the two sets: $C_{ij} = C_{i+\frac{N}{2}\,j} = C_{i\,j+\frac{N}{2}} = C_{i+\frac{N}{2}\,j+\frac{N}{2}}$ for $i, j = 1, \ldots, N/2$. This can be the consequence of fully correlated fluctuations, $C_{ij} = C\ \forall\ i, j$. Alternatively, such a situation arises for example for atoms initially arranged symmetrically with respect to an anti-node such that $G_1^{(0)} = G_2^{(0)}$, if the two atoms (or ions) in each pair l ($l = 1, \ldots, N/2$) are locked into a common oscillation. This should be the case for two trapped ions repelling each other through a strong Coulomb interaction, and cooled below the temperature corresponding to the frequency of the breathing mode. Equation (S5) then gives $\delta\alpha_f = 0$, i.e. no additional noise from the fluctuations of the couplings. Note, however, that for initial $G_1^{(0)} \neq G_2^{(0)}$ the more general DFS leads to a more complicated condition for the correlations, $C_{ij}|G_2^{(0)}|^2 + C_{i+\frac{N}{2}\,j+\frac{N}{2}}|G_1^{(0)}|^2 - C_{i\,j+\frac{N}{2}}G_2^{(0)}G_1^{(0)} - C_{i+\frac{N}{2}\,j}G_1^{(0)}G_2^{(0)} = 0$, which might be harder to achieve.

3. Correlated fluctuations within a set, but uncorrelated between the two sets, $C_{ij} = C$ for $i, j \in \{1, \ldots, N/2\}$ or $i, j \in \{N/2+1, \ldots, N\}$, but $C_{ij} = 0$ for $i \in \{1, \ldots, N/2\}$ and $j \in \{N/2+1, \ldots, N\}$ or vice versa. This case leads to a noise of order $\mathcal{O}(N^2)$, the worst case scenario. However, this comes as no surprise, as such correlations are indistinguishable from the signal: all the atoms in a given set move in a correlated fashion, but independently from the atoms of the other set. This modifies the couplings in the same fashion as if the length of the cavity was changed.

Case (2) above is clearly the most favorable situation. If there are no other background signals depending on N, we keep the $1/N$ scaling of δx. In order to favor case (2) over cases (1),(3), it appears to be advantageous to work with ions and to try to bring the ions in a pair as closely together as possible, thus strongly correlating their fluctuations, while separating the ions in the same set as far as possible.

Imperfections in initial state preparation

Suppose that instead of the state $|\psi_0\rangle$, the state of the atoms

$$|\tilde{\psi_0}\rangle = \bigotimes_{l=1}^{N/2}|\varphi\rangle_l \text{ with } |\varphi\rangle_l = a|t_-\rangle_l + b|s\rangle_l + c|t_0\rangle_l + d|t_+\rangle_l \tag{S6}$$

was prepared (we consider the same state for all pairs for simplicity, but this is not essential, and assume the state normalized). Then,

$$\begin{aligned}\alpha \;=\; \gamma\Big[&\frac{N}{2}\Big(\big((c-b)G_1 + (c+b)G_2\big)^2 + 2d^2(G_1^2+G_2^2)\Big)\\ &+\frac{N}{2}\left(\frac{N}{2}-1\right)\Big((G_2-G_1)b(a-d) + (G_2+G_1)c(a+d)\Big)^2\Big]/g^2\,.\end{aligned} \tag{S7}$$

The derivative of α with respect to G_2 is of order $\mathcal{O}(N^2)$, and thus still allows to find the minimum of α as function of the position of the second lattice with a precision of order $\mathcal{O}(1/N)$. At the minimum a component outside the DFS persists, such that photons will leak out of the cavity, but the average rate is only of order $\mathcal{O}(N)$, and can be measured separately and subtracted from the signal. Changes δG_2 of G_2 away from the position of the minimum still lead to a signal that scales, for large N as N^2, $\alpha = \frac{\gamma}{4}\left(a(c+b)+d(c-b)\right)^2 N^2 \delta G_2^2/g^2$, and the analysis leading to the $1/N$ scaling of δx still applies.

Alternatively, one can get rid of the additional background by letting the system relax before measuring changes of L. Indeed, any state with a component outside the DFS will relax to a DFS state or mixtures of DFS states within a time of order $1/\gamma$ or less. Components with large total pseudo-angular momentum J relax in fact in much shorter time of order $1/(J\gamma)$. The DFS states reached through relaxation starting from $|\tilde{\psi}_0\rangle$ still allow a scaling of α close to N^2. We have shown this by simulating the relaxation process with the help of the stochastic Schrödinger equation (24). Using an Euler scheme with a time step of $0.01/\gamma$, we followed the convergence of $\psi(t)$ to DFS states for states with $(a,b,c,d) = (\cos\delta, 1, 0, \sin\delta)/\sqrt{2}$, until the norm of the difference $|\psi(t+dt)\rangle - |\psi(t)\rangle$ dropped below 10^{-12}. In these final dark states, randomly distributed over the DFS, we calculated α, and averaged over a large number n_r of realizations of the stochastic process ($n_r = 10^5$, 10^4, 10^4, $2.5\cdot10^3$, 10^3, $1.25\cdot10^3$, 250, 200, and 250 for $N = 2,4,6,8,10,12,14,16,$ and 18). Supplementary Figure S1 shows the scaling of α as function of N for different values of δ for $0 \le \delta \le \pi/2$ up to $N = 18$. Within this numerically accessible range of N, α follows a power law $\alpha \propto N^p$ with an exponent p that decays only gradually with δ for $\delta \le \pi/4$. Moreover, that decay might be a finite size effect: Note that, surprisingly, $\alpha(\delta)$ appears to be close to symmetric with respect to $\delta = \pi/4$. This is corroborated by exact analytical calculations based on the diagonalisation of $L_I(x)$, which lead to $\alpha = 1/2$ for $N = 2$, $\alpha = (55 - 12\sin(2\delta) - \cos(4\delta))/36$ for $N = 4$, and $\alpha = (303 - 110\sin(2\delta) - 3\cos(4\delta))/100$ for $N = 6$ (in units $\gamma|G_1 - G_2|^2/g^2$). The plot shows that all numerical data can be very well fitted by $\alpha = A + B\sin(2\delta) + C\cos(4\delta)$. From eq. (21) we know that $A+C$ has to scale as N^2 for sufficiently large N. Both B and C are negative for all N for which we have data, and C appears to be negligible. Supplementary Figure S1 shows that $-B$ increases even more rapidly than N^2 (a fit in the range $N = 8,\ldots,18$ gives a power law $N^{2.4}$). But B has to cross over to a power law N^p with $p \le 2$, unless other Fourier components start contributing significantly. Otherwise, α would become negative for $\delta > 0$. This indicates that for large N the scaling of α is in fact N^2 for all δ.

In summary, our method still works, even if the product state $|\psi_0\rangle$ is not prepared perfectly. One has the choice to start measurement immediately after state preparation, which gives an additional background of order N, or to wait a time of the order of a few $1/\gamma$ after preparation of the initial state, until no more photons leave the cavity through the mirror, with no additional background penalty. In both cases the scaling of the uncertainty of $\delta L/L$ in a subsequent measurement of a small change of L is still $\propto 1/N$.



Supplementary Figure S1: Effects of imperfect state preparation. **(a)** Scaling of collective photon escape rate $\alpha = -\langle \dot{J}_z(x,0)\rangle$ (in units $\gamma|G_1 - G_2|^2/g^2$) in the mixture of DFS states reached by relaxation from state $|\tilde{\psi}_0\rangle$, eq. (S6). Data for $\delta = \delta_0$ ($\delta = \pi/2 - \delta_0$) denoted by squares and full lines (diamonds and dashed lines), respectively; $\delta_0 = 0, \pi/9, 2\pi/9, 3\pi/9$ and $4\pi/9$ in black, red, green, blue, and brown. Exact analytical results for $\delta = 0$, eq. (21) shown with black crosses. Full and dashed lines are guides to the eye only. **(b)** Dependence of α on imperfection parameter δ. $N = 2$, 4,6,8,10,12,14,16,18 in black circles, red squares, green diamonds, blue triangles up, brown triangles left, grey triangles down, violet triangles right, cyan pluses, orange Xs, respectively. The full lines for N=2,4,6 are exact analytical results. The dashed lines for $N = 8,\ldots,18$ are fits to $A + B\sin(2\delta) + C\cos(4\delta)$. The inset shows the scaling of the coefficients A (black circles), $-B$ (red squares), and $-C$ (green diamonds) as function of N.

Supplementary References

[43] Bloch, I., Quantum coherence and entanglement with ultracold atoms in optical lattices, *Nature* **453**, 1016–1022 (2008).